

COHO PRE-SPAWN MORTALITY ON LONGFELLOW CREEK



2016

Survey Results

The Longfellow Creek Watershed drains over 2,000 acres of West Seattle and suffers from urban stream syndrome. Unidentified chemical contaminants in the stormwater runoff that flows into Longfellow Creek negatively impact the physiology of coho salmon, and affected individuals exhibit a suite of symptoms including disorientation, loss of swimming ability, gaping of the mouth, and usually death within hours of symptom onset. Prior surveys of Longfellow Creek reported abnormally high levels of pre-spawn mortality (PSM) in female coho. To continue to document this phenomenon, volunteers trained by Puget Soundkeeper performed daily surveys on Longfellow Creek from October to December 2016, recording sightings of live coho salmon and noting their behavior. Volunteers also performed necropsies on 50 coho carcasses, noting the spawning condition of females. PSM was 48% for the 2016 run.

Coho Pre-spawn Mortality on Longfellow Creek

SUMMARY OF 2016 SURVEYS

BACKGROUND

Ongoing physical and biological restoration efforts sponsored by the City of Seattle and others began on Longfellow Creek in West Seattle, WA in the 1990s (Scholz et al. 2011). Longfellow Creek drains 2,685 acres of West Seattle (King County 2000) and suffers from “urban stream syndrome,” a condition marked by frequent flooding after rainfall events, channelization, and reduced species richness and diversity (Meyer et al. 2005). From 2002 to 2009, various federal and state agencies conducted surveys on coho (*Oncorhynchus kisutch*), Chinook (*O. tshawytscha*), and chum (*O. keta*) salmon that use Longfellow Creek as well as other urban creeks in Puget Sound as spawning habitat to gauge the effectiveness of stream restoration efforts in areas throughout the Puget Sound basin (Scholz et al. 2011). They noticed coho salmon uniquely suffered from exposure to urban stormwater runoff, but Chinook and chum did not appear to experience the same distress (Scholz et al. 2011). When public funding was no longer available to continue these survey efforts, Puget Soundkeeper trained community volunteers to collect salmon run data and continue to monitor trends in spawning success and salmon survival. Volunteer surveys began in 2015 and will continue until agency funding resumes.



Longfellow Creek (A) at base level flow and (B) in flooded conditions.

Urban Salmon Migrations

During the salmon run each fall, a population of coho salmon returns from the open ocean to Elliott Bay, then enters the Duwamish River. After adjusting to a lower salinity, some of these salmon then swim through a grated culvert south of Terminal 5 into the mouth of Longfellow Creek in search of spawning habitat. From there, the salmon follow an underground pipeline for approximately 800 meters before daylighting just south of SW Yancy Street in West Seattle.

Wild salmon migrate to their natal streams using currents and chemical cues (Johnsen and Hasler 1980). Most of the coho that migrate to Longfellow Creek, however, are from hatchery stock and do not have a natal stream to return to, so they navigate to Longfellow as one of the first freshwater inputs they encounter.

Stormwater Toxicity and Coho Pre-spawn Mortality

Migrating salmon in urban waterways like Longfellow Creek are exposed to a chemical cocktail of pollutants flushed into the waterway via stormwater runoff, including petroleum products, rubber from tires, copper from automotive brake pads, combustion products from engines, fertilizers, and pathogens from dog feces and sewage. This mixture of contaminants causes a unique set of symptoms in coho, including disorientation, loss of swimming ability, gaping of the mouth, and usually death within hours of exposure (Scholz et al. 2011). To see an example of these symptoms in Longfellow Creek, follow the link <https://vimeo.com/111234620>.



A coho salmon in Longfellow Creek exhibiting signs of stress.

Coho salmon affected by urban stormwater runoff often die before releasing their eggs or sperm, resulting in a phenomenon called pre-spawn mortality (PSM), which is alarmingly high in urban waterways like Longfellow Creek. In surveys conducted from 2002 to 2009, PSM in female coho was 67 to 100% in Longfellow Creek, compared to 1% in a non-urban stream (Scholz et al. 2011).

PSM in coho was historically 67 to 100% in Longfellow, compared to 1% in a non-urban stream.

Collaborative research efforts conducted by Dr. Jenifer McIntyre at Washington State University, Washington's Department of Fish and Wildlife, and NOAA suggest the contaminants in urban stormwater

runoff are lethal to juvenile coho (McIntyre et al. 2015) as well as adult spawners (Scholz et al. 2011; Spromberg et al. 2016; Washington State University 2016). Exposure to stormwater runoff during coho salmon development has also been linked to an inability to hatch or delayed hatching, smaller body size, atypical bleeding in the brain, smaller eye size, accumulation of fluid around the heart, and deformities in the head and the heart (Washington State University 2016).

Blood analyses of coho salmon exposed to urban stormwater runoff suggest the toxicity causes hypoxia, a condition where the fish are unable to use the oxygen in their tissues for respiration (Washington State University 2016). Interestingly, the level of dissolved oxygen in the water in Longfellow Creek is normal, but the coho are unable to use this oxygen in respiration (Washington State University 2016).

SURVEY PROCEDURE AND RESULTS


Volunteers trained by Puget Soundkeeper using the protocol created by NOAA's Northwest Fisheries Science Center and the Department of Fish and Wildlife surveyed a quarter mile stretch of Longfellow Creek, beginning east of the Dragonfly Pavilion on 28th Avenue SW and SW Dakota Street, Seattle, WA, 98116 and ending just before the bridge on SW Genesee Street and the West Seattle Golf Course. Daily surveys began on October 10th and finished on December 15th, 2016.



King 5 News filming salmon survey volunteers as they examine a deceased female coho.

Volunteers recorded sightings of live fish, noting their species and behavior. They also recorded whether or not each individual fish had an adipose fin, a small fin located behind the dorsal fin. If the fin was clipped, the fish was assumed to be from hatchery stock. Volunteers also performed necropsies on dead coho found in the creek, which included measuring, dissecting, and evaluating the spawning condition of each salmon.

Necropsy Procedure



Necropsy measurements recorded for each deceased coho.

The various measurements taken for each coho carcass included the fork length (length from the tip of the jaws to the end of the tail), the post-orbital to hypural plate length (POH, from behind the coho's eye to the point at which the tail bends), and the girth (circumference just in front of the dorsal fin). The abdomen of each coho was then slit from the vent to the ventral fin, and the body cavity examined to determine spawning condition. Male spawning condition was always marked as unknown, as males can spawn multiple times in a salmon run, and there was no visible evidence whether or not the male coho released their sperm. Female coho were marked as pre-spawn mortality cases (PSM) if $\geq 50\%$ of eggs remained after death. If the female's body cavity was empty or contained $< 50\%$ eggs, she was marked as a post-spawn mortality (POST) case. Any individuals that showed signs of predation were marked as having an unknown (UNK) spawning condition at the time of death.



A female coho marked as a pre-spawn mortality (PSM) case.

2016 Survey Results

Volunteers conducted 60 daily surveys from October 10th to December 15th, 2016, weather permitting. Volunteers recorded 113 sightings of live coho, only one of which was potentially symptomatic. PSM was 48% in female coho.

Sex of Coho	Necropsy Totals	Spawning Condition			
		PSM	% PSM	POST	UNK
Female	21	10	48	11	0
Male	28	---	---	---	28
Unknown	1	---	---	---	1
TOTALS	50	10	48	11	29

For the 50 necropsied fish, the average fork length was 60cm, average POH length was 44cm, and average girth was 30cm. All necropsied coho were from hatchery stock. The coho salmon necropsied this year had a larger average body size than those dissected in 2015, with a 16cm longer average fork length and 6cm larger average girth.

CONCLUSIONS

The number of coho recorded in Longfellow Creek in 2016 was encouraging, especially after the coho salmon run was predicted to be lower than normal and fishing for this species was initially restricted. These numbers contrast with results gathered by Puget Soundkeeper and volunteers in 2015, with only 31 necropsies performed on coho in that year (22 male, 9 female with 1 PSM mortality case). Additionally, most of the males necropsied in 2015 were “jack” (juvenile) males, whereas only one jack male was recorded in 2016, likely explaining the larger average body size of the coho necropsied in 2016 versus 2015.

A pre-spawn mortality level of 48% in female coho was lower than that seen in previous years, but it would be premature to suggest any sort of permanent downward trend in PSM in Longfellow Creek. Interestingly, rainfall was 12.3cm above average in October through December for this year with an especially rainy October. Increased rainfall has been qualitatively correlated with decreased PSM on Longfellow Creek (Scholz et al. 2011), perhaps because the chemical contaminants in stormwater are more frequently flushed through the watershed and do not have the chance to accumulate to levels that are toxic to coho. Additionally, ongoing restoration efforts at Longfellow Creek continue to improve the local ecology of this urban ecosystem and may also have contributed to a decreased toxicity in the area’s stormwater runoff during the survey season. For example, Delridge Neighborhood Development Association’s Wetlands and Stewardship Project constructed a 1 acre wetland to help filter runoff before it flows into Longfellow Creek. The City of Seattle has developed a number of programs and policies aimed at filtering and reducing stormwater flow to streams. The Nature Consortium also works with volunteers to remove invasive species like Himalayan blackberry from the banks of the creek and replace them with native, shade-providing species. The collective efforts of these groups and others to improve Longfellow Creek may also be partially responsible for the decreased level of PSM and reduced toxicity symptoms observed in coho in 2016.

Past research also suggests that installing simple systems to filter urban stormwater runoff can have a significant impact on coho survival (McIntyre et al. 2015; Spromberg et al. 2016; Washington State University 2016). In laboratory studies, filtering urban stormwater runoff through a mixture of 60% sand and 40% compost with a topping of mulched bark reduced coho salmon mortality from 100% to 0% in both adults (Spromberg et al. 2016) and juveniles (McIntyre et al. 2015). Adopting similar filtration methods on a larger scale, called Green Stormwater Infrastructure (GSI), includes systems like rain gardens, swales, and permeable pavement. Installing GSI on a biologically relevant scale in the Longfellow Creek Watershed has the potential to drastically reduce coho PSM.

This research provides an example of community based education and outreach, as well as a visually impactful way to spread awareness around urban stormwater runoff and its ecological effects. Puget Soundkeeper would not have been able to gather these data without the help of dedicated volunteers, who committed their time and resources every week to document the plight of coho salmon in this urban environment.



A salmon survey volunteer analyzing a coho carcass.

REFERENCES

- Johnsen, PB, and AD Hasler. 1980. "The Use of Chemical Cues in the Upstream Migration of Coho Salmon, *Oncorhynchus kisutch* Walbaum." *Journal of Fish Biology* 17 (1): 67-73. doi:10.1111/j.1095-8649.1980.tb02742.x.
- King County. Department of Natural Resources and Parks, Water and Land Resources Division. 2000. *WRIA 9 Habitat Limiting Factors and Reconnaissance Assessment for Salmon Habitat Part 11: Factors of Decline/Conditions 3.1 Longfellow Creek Subbasin*. <http://www.govlink.org/watersheds/9/reports/Recon.aspx>.
- McIntyre, JK, Davis, JW, Hinman, C, Macneale, KH, Anulacion, BF, Scholz, NL, and JD Stark. 2015. "Soil Bioretention Protects Juvenile Salmon and Their Prey from the Toxic Impacts of Urban Stormwater Runoff." *Chemosphere* 132: 213-219. doi:10.1016/j.chemosphere.2014.12.052.
- Meyer, Judy L., Paul, Michael J., and W. Keith Taulbee. 2005. "Stream Ecosystem Function in Urbanizing Landscapes." *Journal of the North American Benthological Society* 24 (3): 602-612. doi:10.1899/04-021.1.
- Scholz, Nathaniel L., Myers, Mark S., McCarthy, Sarah G., Labenia, Jana S., McIntyre, Jenifer K., Ylitalo, Gina M., Rhodes, Linda D. et al. 2011. "Recurrent Die-Offs of Adult Coho Salmon Returning to Spawn in Puget Sound Lowland Urban Streams." *PLOS ONE* 6 (12): 1-12. doi:10.1371/journal.pone.0028013.
- Spromberg, Julann A., Baldwin, David H., Damm, Steven E., McIntyre, Jenifer K., Huff, Michael, Sloan, Catherine A., Anulacion, Bernadita F., Davis, Jay W., and Nathaniel L. Scholz. 2016. "Coho Salmon Spawner Mortality in Western US Urban Watersheds: Bioinfiltration Prevents Lethal Storm Water Impacts." *Journal of Applied Ecology* 53 (2): 398-407. doi:10.1111/1365-2664.12534.
- Washington State University. "WSU Innovators Event: Dr. Jenifer McIntyre." Filmed [April 2016]. YouTube video, 33:04. Posted [April 2016]. <https://www.youtube.com/watch?v=BNruLbD2sP8>.